

---

**CIRM Marseille 15-19 January 2018**

**Kähler immersions of homogeneous Kähler manifolds  
into complex space forms**

**Andrea Loi, University of Cagliari**

**joint works (2006-2018) with:**

**A. J. Di Scala, H. Hishi, R. Mossa, F. Salis, M. Zedda, F. Zuddas**

---

**Aim** *Classify all homogeneous Kähler manifolds (h.K.m.) which admit a Kähler immersion into a given finite or infinite dimensional complex space form.*

**Advertising for the book:** -, M. Zedda, *Kähler immersions of Kähler manifolds into complex space forms*, <https://arxiv.org/abs/1712.>

---

1. General definitions: Kähler manifolds, Complex space forms and their classifications.

2. Kähler immersions into complex space forms, E. Calabi, Ann. Math. 1953.

3. Homogeneous Kähler manifolds and their classification (J. Dorfmeister, K. Nakajima, Acta Math. 1988).

4. Kähler immersions of h.K.m. into complex space forms (Theorem 1, 2, 3, 4).

5. Sketch of the proofs of Theorem 1, 2, 3, 4.

6. The Kähler-Einstein case

.

---

**1. General definitions: Kähler manifolds,  
complex space forms and their classification**

---

## Kähler manifolds

Let  $(M, g) = (M, g, \omega, J)$  be a Kähler manifold of complex dimension  $n$ .

$$\omega(X, Y) = g(X, JY), \quad X, Y \in \mathfrak{X}(M), \quad d\omega = 0.$$

The form  $\omega$  is called the *Kähler form associated to the metric  $g$* .

On a contractible open set  $U \subset M$

$$\omega = \frac{i}{2} \partial \bar{\partial} \Phi = \frac{i}{2} \sum_{j,k=1}^n \frac{\partial^2 \Phi}{\partial z_j \partial \bar{z}_k} dz_j \wedge d\bar{z}_k,$$

where  $\Phi : U \rightarrow \mathbb{R}$  is a strictly PSH function called a *Kähler potential* for the metric  $g$ .

---

## Complex space forms

A *complex space form*  $(S, g_S) = (S, g_S, \omega_S, J_S)$  is a finite or infinite dimensional Kähler manifold of constant holomorphic sectional curvature.

---

## Classification of complex space forms

Complex Euclidean space  $\mathbb{C}^{N \leq \infty} := (\mathbb{C}^{N \leq \infty}, g_0)$

$\mathbb{C}^\infty := \ell^2(\mathbb{C})$  ( $z = \{z_j\} \in \ell^2(\mathbb{C})$  iff  $\sum_{j=1}^\infty |z_j|^2 < \infty$ )

$$\omega_0 = \frac{i}{2} \partial \bar{\partial} |z|^2 = \frac{i}{2} \sum_{j=1}^N dz_j \wedge d\bar{z}_j, \quad |z|^2 = |z_1|^2 + \cdots + |z_N|^2.$$

Complex hyperbolic space  $\mathbb{C}H^{N \leq \infty} := (\{z \in \mathbb{C}^N \mid |z|^2 < 1\}, g_{hyp})$

$$\omega_{hyp} = -\frac{i}{2} \partial \bar{\partial} \log(1 - |z|^2).$$

Complex projective space  $\mathbb{C}P^{N \leq \infty} = (\mathbb{C}^{N+1} \setminus \{0\} / z \sim \lambda z, g_{FS})$

$$\omega_{FS}|_{U_0} = \frac{i}{2} \partial \bar{\partial} \log(1 + |z|^2), \quad z_j = \frac{Z_j}{Z_0}, \quad j = 1, \dots, N, \quad U_0 = \{Z_0 \neq 0\}.$$

---

**2. Kähler immersions into complex space forms  
(E. Calabi, Ann. Math. 1953)**



---

## Kähler immersions into complex space forms

Let  $(M, g)$  be a Kähler manifold. A Kähler immersion

$$f : (M, g) \rightarrow (S, g_S)$$

is a holomorphic map (i.e.  $df \circ J = J_S \circ df$ ) which is isometric (i.e.  $f^*g_S = g$ ).

**Remark** The “starting” manifold  $M$  will be always finite dimensional.

**Terminology:** A Kähler metric  $g$  on a complex manifold  $M$  is *projectively induced* if  $(M, g)$  can be Kähler immersed into a finite or infinite dimensional complex projective space.

---

**Calabi's results on Kähler immersions (1953)**

**Theorem** (Calabi's rigidity) *Let  $f : (M, g) \rightarrow (S, g_S)$  be a Kähler immersion. Then any other Kähler immersion of  $(M, g)$  into  $(S, g_S)$  is given by  $\mathcal{U} \circ f$  where  $\mathcal{U}$  is a unitary transformation, i.e.  $\mathcal{U} \in \text{Aut}(S) \cap \text{Isom}(S, g_S)$ .*

**Theorem** (Calabi's extension theorem) *A simply-connected Kähler manifold  $(M, g)$  admits a Kähler immersion into a given complex space form  $(S, g_S)$  iff there exists an open set  $U \subset M$  such that  $(U, g|_U)$  can be Kähler immersed into  $(S, g_S)$ .*

---

## Complex Euclidean spaces into complex space forms

$$\mathbb{C}^n \rightarrow \mathbb{C}H^{N \leq \infty}, \mathbb{C}P^{N < \infty}$$

$$\mathbb{C}^n \hookrightarrow \mathbb{C}^{N \leq \infty}, n \leq N$$

We have the following **Calabi's immersion**

$$\mathbb{C}^n \rightarrow \mathbb{C}P^\infty : z \mapsto (\dots, \sqrt{\frac{1}{j!}} z^j, \dots), |j| \geq 0$$

$$z^j = z_1^{j_1} \cdots z_n^{j_n} \quad |j| = j_1 + \cdots + j_n, \quad j! = j_1! \cdots j_n!$$

---

## Complex hyperbolic spaces into complex space forms

Let  $\mathbb{C}H_\lambda^n = (\mathbb{C}H^n, \lambda g_{hyp})$ ,  $\lambda > 0$ ,  $\mathbb{C}H^n := \mathbb{C}H_1^n = (\mathbb{C}H^n, g_{hyp})$

$\mathbb{C}H_\lambda^n \rightarrow \mathbb{C}^{N < \infty}, \mathbb{C}P^{N < \infty}$

$\mathbb{C}H_\lambda^n \rightarrow \mathbb{C}H^{N \leq \infty} \Leftrightarrow \lambda = 1, n \leq N$

We have the following [Calabi's immersions](#)

$$\mathbb{C}H_\lambda^n \rightarrow \ell^2(\mathbb{C}) : z \mapsto \sqrt{\lambda}(\dots, \sqrt{\frac{(|j| - 1)!}{j!}} z^j, \dots), \quad |j| \geq 1$$

$$\mathbb{C}H_\lambda^n \rightarrow \mathbb{C}P^\infty : z \mapsto (\dots, \sqrt{\frac{\lambda(\lambda + 1) \cdots (\lambda - 1 + |j|)}{j!}} z^j, \dots), \quad |j| \geq 0$$

$$z^j = z_1^{j_1} \cdots z_n^{j_n}, \quad |j| = j_1 + \cdots + j_n, \quad j! = j_1! \cdots j_n!$$

---

## Complex projective spaces into complex space forms

Let  $\mathbb{C}P_\lambda^n = (\mathbb{C}P^n, \lambda g_{FS})$ ,  $\lambda > 0$ ,  $\mathbb{C}P^n := \mathbb{C}P_1^n = (\mathbb{C}P^n, g_{FS})$

$$\mathbb{C}P_\lambda^n \rightarrow \mathbb{C}^{N \leq \infty}, \mathbb{C}H^{N \leq \infty}$$

Let  $k \in \mathbb{Z}$  and  $N_k := \frac{(n+k)!}{n!k!} - 1$ . Then the map

$$\mathbb{C}P_k^n \xrightarrow{V_k} \mathbb{C}P^{N_k} : [Z] \mapsto [\dots, \sqrt{\frac{|j|!}{j!}} Z^j, \dots], \quad |j| \geq 0$$

$Z^j = Z_0^{j_0} \cdots Z_n^{j_n}$ ,  $|j| = j_0 + \cdots + j_n$ ,  $j! = j_0! \cdots j_n!$  satisfies

$$V_k^* g_{FS} = k g_{FS}$$

---

**3. H.K.m. and their classification  
(J. Dorfmeister, K. Nakajima, 1988)**

---

## Homogeneous Kähler manifolds

*A homogeneous Kähler manifold (h.K.m.) is a Kähler manifold  $(M, g)$  such that the Lie group  $G = \text{Aut}(M) \cap \text{Isom}(M, g)$  acts transitively on  $M$ .*

**Remark.** The metric  $g$  is not uniquely determined by  $G$ . There exist different (neither homothetic or isometric)  $G$ -invariant homogeneous metrics.

**Examples:** Complex space forms are h.K.m.

---

## Homogeneous bounded domains

Let  $\Omega \subset \mathbb{C}^n$ ,  $\Omega$  bounded domain endowed with a homogeneous Kähler metric  $g_\Omega$ . Then  $(\Omega, g_\Omega)$  is called a *homogeneous bounded domain* (h.b.d.).

If  $\text{Aut}(\Omega)$  acts transitively on  $\Omega \subset \mathbb{C}^n$  then  $(\Omega, g_\Omega = g_B)$  is a bounded symmetric domain \*,  $g_B$  is the Bergman metric whose associated Kähler form  $\omega_B = \frac{i}{2} \partial \bar{\partial} \log K$ , where  $K$  is the reproducing kernel for the Hilbert space of holomorphic  $L^2$ -functions on  $\Omega$ .

**Remark.** Every bounded symmetric domain  $(\Omega, g_B)$  is a h.b.d. but there exist (Pyatetskii-Shapiro, 1969) h.b.d.  $(\Omega, g_B)$  which are not bounded symmetric domains.

\*A bounded symmetric domain  $\Omega \subset \mathbb{C}^n$  is a domain where the geodesic symmetry  $\exp_x(v) \mapsto \exp_x(-v)$  is a Kähler map.



---

**Other examples of h.K.m.**

Flat h.K.m.  $\mathcal{E} = \mathbb{C}^k \times F$  where  $F$  is a non simply-connected (either compact or non compact) flat Kähler manifold.

Compact simply-connected h.K.m. These are also called *Kähler C-spaces* or *rational homogeneous varieties*.

Compact h.K.m.  $(M, g) = \mathcal{C} \times T_1 \times \cdots \times T_l$ ,  $\mathcal{C}$ -space,  $T_j$  flat torus.

Products of homogeneous Kähler manifolds The products of h.K.m. is a h.K.m.

---

**Solution of the fundamental conjecture (FC) for h.K.m.**

**Theorem FC** (J. Dorfmeister, K. Nakajima, 1988) *A h.K.m.  $(M, g)$  is the total space of a holomorphic fiber bundle over a h.b.d.  $(\Omega, g_\Omega)$ . Moreover the fiber  $\mathcal{F} = \mathcal{E} \times \mathcal{C}$  is (with the induced Kähler metric) the Kähler product of a flat homogeneous Kähler manifold  $\mathcal{E} = \mathbb{C}^k \times F$  and a  $\mathcal{C}$ -space  $\mathcal{C}$ .*

$$\begin{array}{ccc} \mathcal{F} = \mathcal{E} \times \mathcal{C} & \xrightarrow{\text{Kähler}} & (M, g) \\ & & \pi \downarrow \\ & & (\Omega, g_\Omega) \end{array}$$

**Remark.**  $M \stackrel{top}{\cong} \Omega \times \mathcal{F}$  as a complex manifold.

---

**4. Kähler immersions of h.K.m.  
into complex space forms (Theorems 1, 2, 3, 4)**

---

**Homogeneous Kähler manifolds into  $\mathbb{C}^{N \leq \infty}$**

**Theorem 1** (-, A. J. Di Scala, H Hishi, 2012) *Let  $(M, g)$  be a  $n$ -dimensional h.K.m. which can be Kähler immersed into  $\mathbb{C}^{N \leq \infty}$ . Then  $(M, g) = \mathbb{C}^k \times \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l}$ . Moreover, the immersion is given, up to a unitary transformation of  $\mathbb{C}^N$  by*

$$f_0 \times f_1 \times \cdots \times f_l,$$

where  $f_0$  is the linear inclusion  $\mathbb{C}^k \xrightarrow{\text{tot.geod.}} \mathbb{C}^N$  and each  $f_r : \mathbb{C}H_{\lambda_r}^{n_r} \rightarrow \ell^2(\mathbb{C})$ ,  $r = 1, \dots, l$ , are Calabi's immersions.

---

**Homogeneous Kähler manifolds into  $\mathbb{C}H^{N \leq \infty}$**

**Theorem 2** (-, A. J. Di Scala, H Hishi, 2012) *Let  $(M, g)$  be a  $n$ -dimensional h.K.m. which can be Kähler immersed into  $\mathbb{C}H^{N \leq \infty}$ . Then, up to a unitary transformation of  $\mathbb{C}H^N$ ,*

$$(M, g) = \mathbb{C}H^n \xrightarrow{\text{tot.geod.}} \mathbb{C}H^N.$$

---

**Two theorems on h.K.m. into  $\mathbb{C}P^{N \leq \infty}$**

**Theorem 3** (-, A. J. Di Scala, H Hishi, 2012) *Let  $(M, g)$  be a  $n$ -dimensional h.K.m. which can be Kähler immersed into  $\mathbb{C}P^{N \leq \infty}$ . Then  $\omega$  is integral,  $\pi_1(M) = 1$  and the immersion is injective.*

**Theorem 4** (-, R. Mossa, 2014) *Let  $(M, g)$  be a **simply-connected** h.K.m. such that its associated Kähler form  $\omega$  is **integral**. Then there exists  $m_0 \in \mathbb{Z}$  such that*

$$(M, m_0 g) \rightarrow \mathbb{C}P^{N \leq \infty}.$$

---

## Remarks on the compact case

When  $M$  is **compact** Theorem 3 and Theorem 4 were proved by M. Takeuchi (1978) using the theory of semisimple Lie groups and Dynkin diagrams (one can take  $m_0 = 1$  in Theorem 4).


Notice that if a h.K.m. can be Kähler immersed into  $\mathbb{C}P^{N < \infty}$  then  $M$  is a  $C$ -space, i.e. is a **compact** (simply-connected) Kähler manifold.

Viceversa if  $M$  is any **compact** (not necessarily homogeneous) Kähler manifold which can be Kähler immersed into  $\mathbb{C}P^{N \leq \infty}$  one can assume  $N < \infty$ .

---

**The necessity of taking  $m_0$  in Theorem 4**

Let  $\Omega$  be an irreducible bounded symmetric domain. The Wallach set<sup>†</sup>  $W(\Omega) \subset \mathbb{R}$  is a subset of  $\mathbb{R}$  which “looks like”:



**Important property of the Wallach set:**  $W(\Omega) = \mathbb{R}$  (and hence the discrete part of  $W(\Omega)$  is empty) if and only if  $\Omega = \mathbb{C}H^n$ .

<sup>†</sup> $W(\Omega)$  consists of all  $\lambda \in \mathbb{R}$  such that there exists a Hilbert space  $\mathcal{H}_\lambda$  whose reproducing kernel is  $K_\lambda^\gamma$ ,  $\gamma$  the genus of  $\Omega$ , where  $K$  is the reproducing kernel for the Hilbert space of holomorphic  $L^2$ -functions on  $\Omega$ .



---

## The Wallach set and Kähler immersions into $\mathbb{C}P^\infty$

**Theorem W** (–, M. Zedda, 2010) *Let  $(\Omega, g_B)$  be a irreducible bounded symmetric domain. Then  $(\Omega, \lambda g_B)$  can be Kähler immersed into  $\mathbb{C}P^\infty$  if and only if  $\lambda\gamma \in W(\Omega) \setminus \{0\}$ , where  $\gamma$  denotes the genus of  $\Omega$ .*

---

## Two consequences of Theorem W

**First consequence:** Let  $(\Omega, g_B) \neq \mathbb{C}H^n$  be a irreducible bounded symmetric domain. One can find  $\lambda > 0$  such that  $\lambda\gamma \notin W(\Omega)$ :

0   .   .   .   .   \*   \_\_\_\_\_

$\uparrow$   
 $\lambda\gamma \notin W(\Omega)$

By Theorem W,  $\lambda g_B$  is not projectively induced and  $\lambda\omega_B$  is integral (this shows the necessity of taking  $m_0 > 1$  in Theorem 4).

**Second consequence:** The complex hyperbolic space is the only irreducible bounded symmetric domain  $(\Omega, g_B)$  where  $\lambda g_B$  is projectively induced, for all  $\lambda > 0$ .

---

## A Lemma for homogeneous bounded domains

**Lemma H** (-, A. J. Di Scala, H Hishi, 2012) *Let  $(\Omega, g_\Omega)$  be a h.b.d. If  $(\Omega, \lambda g_\Omega)$  can be Kähler immersed into  $\mathbb{C}P^\infty$  for all  $\lambda > 0$ , then  $(\Omega, g_\Omega) = \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l}$ .*

**Ingredients for the proof.** Unitary representation of semisimple Lie groups; reproducing kernels of weighted Bergman spaces.

---

**5. Sketch of the proofs of Theorem 1, 2, 3, 4**

---

**Sketch of the proof of Theorem 1**

$(M, g) \xrightarrow{f} \mathbb{C}^{N \leq \infty}$  we want to prove that:

$$(M, g) = \mathbb{C}^k \times \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l} \text{ and } f = f_0 \times f_1 \times \cdots \times f_l.$$

1. Theorem FC + Calabi's rigidity theorem + max principle  $\Rightarrow$

$$\mathcal{F} = \mathbb{C}^k \times \mathcal{F} \times \mathcal{C} \xrightarrow{\text{Kähler}} (M, g) \rightarrow \mathbb{C}^{N \leq \infty}$$

$$\pi \downarrow$$

$$(\Omega, g_\Omega)$$

2. Riemannian geometry + homogeneity  $\Rightarrow$

$$(M, g) \stackrel{\text{Kähler}}{=} \mathbb{C}^k \times (\Omega, g_\Omega) \Rightarrow (\Omega, \lambda g_\Omega) \rightarrow \mathbb{C}^{N \leq \infty}, \forall \lambda > 0.$$

---

3. S. Bochner (1947)  $\Rightarrow (\Omega, \lambda g_\Omega) \rightarrow \mathbb{C}P^\infty, \forall \lambda > 0.$

4. Lemma H  $\Rightarrow (\Omega, g_\Omega) = \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l} \Rightarrow$

$\Rightarrow (M, g) = \mathbb{C}^k \times \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l}.$

5. The fact that the immersion  $f$  is, up to a unitary transformation of  $\mathbb{C}^N$ , of the form  $f = f_0 \times f_1 \times \cdots \times f_l$  follows by the reducibility of a Kähler product into  $\mathbb{C}^{N \leq \infty}$  and by Calabi's rigidity theorem.  $\square$

---

**Sketch of the proof of Theorem 2 (based on Theorem 1)**

If  $(M, g) \rightarrow \mathbb{C}H^{N \leq \infty}$  we want to prove that

$$(M, g) = \mathbb{C}H^n \xrightarrow{\text{tot. geod.}} \mathbb{C}H^N.$$

1.  $(M, g) \rightarrow \mathbb{C}H^{N \leq \infty} \Rightarrow (M, g) \rightarrow \ell^2(\mathbb{C})$ .
2. **Theorem 1**  $\Rightarrow (M, g) = \mathbb{C}^k \times \mathbb{C}H_{\lambda_1}^{n_1} \times \cdots \times \mathbb{C}H_{\lambda_l}^{n_l} \Rightarrow M = \mathbb{C}H^n$ .  $\square$

---

**Sketch of the proof of Theorem 3**

Let  $f : (M, g) \rightarrow \mathbb{C}P^{N \leq \infty}$  be a Kähler immersion.

The **integrality** of  $\omega = f^*\omega_{FS}$  is immediate since  $\omega_{FS}$  is integral.

$$\text{Th. FC} \Rightarrow \mathcal{F} = \mathbb{C}^k \times \mathbb{F} \times C \xrightarrow{\text{Kähler}} (M, g) \rightarrow \mathbb{C}P^{N \leq \infty} \Rightarrow M \stackrel{\text{top}}{=} (\Omega, g_\Omega)$$

$\pi \downarrow$

$\Omega \times \mathbb{C}^n \times C$  is **simply-connected**.

Calabi's rigidity  $\Rightarrow f \circ g = \mathcal{U}_g \circ f, \forall g \in G = \text{Aut}(M) \cap \text{Isom}(M, g)$   
 $\Rightarrow f(M)$  is a h.K.m.  $\Rightarrow f(M) \subset \mathbb{C}P^N$  is simply-connected.

$f : M \rightarrow f(M)$  is a local isometry  $\Rightarrow f$  is a covering map  $\Rightarrow f$  is **injective**. □



---

## Sketch of the proof of **Theorem 4**

Let  $(M, g)$  be a **simply-connected** h.K.m. with  $\omega$  **integral** we want to show that  $(M, m_0 g) \rightarrow \mathbb{C}P^{N \leq \infty}$ , for some  $m_0 \in \mathbb{Z}$ .

1. Let  $L$  be a holomorphic line bundle with  $c_1(L) = [\omega]$  and consider the Hilbert space

$$\mathcal{H}_m = \left\{ s \in H^0(L) \mid \int_M h_m(s, s) \frac{\omega^n}{n!} < \infty \right\}$$

where  $h_m$  is an Hermitian metric on  $L^m$  such that  $\text{Ric}(h_m)^\ddagger = m\omega$ .

2. There exists  $m_0 \in \mathbb{Z}$  such that  $\mathcal{H}_{m_0} \neq \{0\}$  (J. Rosenberg, M. Vergne, 1984);

$\ddagger \text{Ric}(h_m) = -\frac{i}{2} \partial \bar{\partial} \log h_m(\sigma(x), \sigma(x))$ , where  $\sigma : U \rightarrow L^m$  is a trivialising holomorphic section of  $L^m$ .

---

3. Consider the smooth function on  $M$  given by:

$$\epsilon_{m_0}(x) = \sum_{j=0}^{d_{m_0}} h_{m_0}(s_j(x), s_j(x)),$$

where  $\{s_0, \dots, s_{d_{m_0}}\}$  is an orthonormal basis of  $\mathcal{H}_{m_0}$ .

**Homogeneity +  $\pi_1(M) = 1 \Rightarrow \epsilon_{m_0}(x)$  is a positive constant.**

4. Therefore the “Kodaira map”

$$\varphi_{m_0} : M \rightarrow \mathbb{C}P^{d_{m_0}}, x \mapsto [s_0(x), \dots, s_{d_{m_0}}(x)]$$

is well-defined and it satisfies

$$\varphi_{m_0}^* \omega_{FS} = m_0 \omega + \frac{i}{2} \partial \bar{\partial} \log \epsilon_{m_0} = m_0 \omega.$$

□

---

## 5. The Kähler-Einstein case

---

## The Kähler-Einstein case

**Theorem** (M. Umehara, 1987) *Let  $(M, g)$  be a complete KE manifold of complex dimension  $n$  which admits a Kähler immersion into  $\mathbb{C}^N$  (resp.  $\mathbb{C}H^N$ ). Then  $(M, g) = \mathbb{C}^n$  (resp.  $(M, g) = \mathbb{C}H^n$ ).*

**Conjecture A:** *A compact KE manifold which admits a Kähler immersion into a complex projective space is homogeneous (Chern (1967), Tsukada (1986), Hulin (2000)).*

**Remark:** *The conjecture cannot be weakened to the noncompact case. There exist examples (even continuous family) of noncompact and nonhomogeneous KE submanifolds of  $\mathbb{C}P^\infty$  (–, M. Zedda, 2010).*

---

## The Ricci flat case

**Conjecture B:** *A Ricci flat projectively induced Kähler metric is flat.*

**Theorem** (-, F. Salis, F. Zuddas, 2018) *A projectively induced Ricci flat and **radial** Kähler metric is flat.*

---

**Conjecture B cannot be weakened to scalar flat metrics**

Let  $S$  be the blow-up of  $\mathbb{C}^2$  at the origin and denote by  $E$  the exceptional divisor. Simanca constructs a scalar flat Kähler complete (not Ricci-flat) metric  $g_S$  on  $S$  whose Kähler potential on  $S \setminus E = \mathbb{C}^2 \setminus \{0\}$  can be written as

$$\Phi_S(|z|^2) = |z|^2 + \log |z|^2, |z|^2 = |z_1|^2 + |z_2|^2.$$

The holomorphic map

$$\varphi : S \setminus E \rightarrow \mathbb{C}P^\infty : (z_1, z_2) \mapsto (z_1, z_2, \dots, \sqrt{\frac{j+k}{j!k!}} z_1^j z_2^k, \dots), j+k \neq 0,$$

is a Kähler immersion. By Calabi's extension theorem it extends to a Kähler immersion of  $(S, g_S)$  into  $\mathbb{C}P^\infty$ .

---

## References

- [1] (joint with M. Zedda) Kähler immersions of Kähler manifolds into complex space forms, preprint 2017, <https://arxiv.org/abs/1712.04298>.
- [2] (joint with Filippo Salis e Fabio Zuddas) Two conjectures on Ricci flat metrics, to appear in Math. Z.
- [3] (joint with Roberto Mossa) Some remarks on Homogeneous Kähler manifolds, *Geom. Dedicata* Volume 179 (2015), 1-7.
- [4] (joint with A. J. Di Scala and H. Hishi) Kähler immersions of homogeneous Kähler manifolds into complex space forms, *Asian J. of Math.* Vol. 16 No. 3 (2012), 479-488.
- [5] (joint with M. Zedda), Kähler-Einstein submanifolds of the infinite dimensional projective space, *Math. Ann.* 350 (2011), 145-154.
- [6] (joint with A. J. Di Scala), Kähler manifolds and their relatives, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) Vol. IX* (2010), 495-501.
- [8] (joint with A.J. Di Scala) Kähler maps of Hermitian symmetric spaces into complex space forms, *Geom. Dedicata* 25 (2007), 103-113.

---

**Thank you for your attention!**